

Estimating Reservoir Competence of *Borrelia burgdorferi* Hosts: Prevalence and Infectivity, Sensitivity, and Specificity

JESSE L. BRUNNER,^{1,2} KATHLEEN LOGIUDICE,³ AND RICHARD S. OSTFELD¹

J. Med. Entomol. 45(1): 139–147 (2008)

ABSTRACT Most vector-borne zoonotic pathogens are transmitted among several host species, but different species vary considerably in their importance to pathogen transmission, at least partially because they vary in their propensity to infect feeding vectors. This propensity is often called realized reservoir competence. Realized reservoir competence is the product of 1) the probability the individual host is infected, i.e., infection prevalence, and 2) the probability that if the host is infected, it will transmit the infection to a feeding vector, or infectivity. Prevalence varies in space and time, whereas infectivity may be a property of the host species. Both prevalence and infectivity are ecologically and epidemiologically important, but measuring them simultaneously is difficult. We present a probabilistic model that separately estimates host infection prevalence and infectivity from data on the infection status of vectors collected from individual hosts, data generally used to measure realized reservoir competence. We then consider how imperfect diagnostic tests (i.e., false negatives and positives) influence these probabilities—estimates of prevalence and infectivity are fairly robust to false negatives, but not to false positives. We thus extend the model to estimate the rate of false positives in order to improve estimates of prevalence and infectivity. We illustrate these methods by reanalyzing data from LoGiudice et al. (2003; Proc. Natl. Acad. Sci. U.S.A. 100: 567–571) on the reservoir competence of ten vertebrate hosts of *Borrelia burgdorferi*, the agent of Lyme disease. We find that these vertebrate hosts vary both in prevalence and infectivity and that both values are highly, positively correlated among species.

KEY WORDS reservoir competence, prevalence, infectivity, *Borrelia burgdorferi*, *Ixodes scapularis*

Most pathogens of animals can infect multiple host species, that is, they are host generalists (Woolhouse et al. 2001). Different host species, however, vary considerably in their importance to pathogen transmission and the dynamics of infection. As a consequence, overall disease incidence—and risk of infection to humans in the case of zoonotic diseases—can be thought of as a function of the host community's composition (Van Buskirk and Ostfeld 1995, Ostfeld and Keesing 2000, Keesing et al. 2006). To understand this relationship and predict actual disease risk, we must be able to measure the contributions of different species to overall rates of transmission.

Mather et al. (1989) introduced the concept of “reservoir potential” as a metric of the contributions of different rodent hosts to transmission of the Lyme disease agent, *Borrelia burgdorferi*, although this concept is equally relevant to other vector-borne zoonoses. Reservoir potential is defined as the average number (or proportion) of infected vectors produced by an individual of a given host species (Fig. 1). It is a product of the number (or propor-

tion) of vectors fed by an individual of a given species and “realized reservoir competence” (Schauber and Ostfeld 2002, LoGiudice et al. 2003)—the probability that a vector feeding on that host species becomes infected. A species with high reservoir potential is one upon which a great number of vectors successfully feed, and from which many become infected with the pathogen. A host species may have low reservoir potential because it feeds few vectors or because it is unlikely to transmit the pathogen to them (low realized reservoir competence), or both. A host species may have low realized reservoir competence, in turn, because individuals are rarely infected (low prevalence) or because when they are infected they are unlikely to transmit the infection to feeding vectors (Fig. 1), a property we call infectivity (“host infectivity” in Kahl et al. 2002).

There are two general approaches to measuring reservoir competence for Lyme disease. In the first approach, an investigator collects host animals from the field and brings them into the laboratory to collect engorged ticks (e.g., LoGiudice et al. 2003) or feeds uninfected xenodiagnostic ticks upon them (e.g., Mather et al. 1989, Levin et al. 1995). These fed ticks are then tested for *B. burgdorferi* infection, usually after molting, to provide a direct measure of realized

¹ Institute of Ecosystem Studies, P.O. Box AB, Millbrook, NY 12545.

² Corresponding author, e-mail: brunnerj@ecostudies.org.

³ Department of Biological Sciences, Union College, Schenectady, NY 12308.

How many infected nymphal ticks are produced by an individual of a given species?




Fig. 1. Relationship among the components of reservoir potential (sensu Mather et al. 1989) and the factors that may influence each. Defined in the context of Lyme disease (see Kahl et al. 2002), these terms assume a permissive vector, such as *Ixodes scapularis* Say.

reservoir competence (Fig. 1). These studies produce an ecologically relevant metric, but they are difficult to generalize from because realized reservoir competence is partially determined by prevalence, which varies in space and time.

The second approach is to first experimentally infect hosts and then, after a given interval, feed xenodiagnostic ticks on these known-infected hosts (e.g., Markowski et al. 1998). These ticks are similarly tested for *B. burgdorferi* infection to provide a direct measure of the probability an infected host will transmit *B. burgdorferi* to a feeding tick (infectivity). Measured values of infectivity reflect host physiology, immunological response(s), and other factors that determine host permissiveness to *B. burgdorferi*. It seems reasonable to assume that infectivity is consistent among members of a species or, in the face of host immune responses, that it at least follows a consistent temporal trend. Intraspecific consistency makes this method useful for comparing the relative *potential* importance of different species to *B. burgdorferi* transmission. However, high estimates of infectivity when hosts are artificially inoculated cannot demonstrate that individuals of a given species play an important role in pathogen transmission in nature. To establish species-specific roles in natural *B. burgdorferi* transmission, one also must estimate natural infection prevalence in each host species.

Unfortunately, the diagnostic tests available for detecting *B. burgdorferi* in vertebrate tissues are not very sensitive, although improving (Eisendle et al. 2007, Wilske et al. 2007) and are thus liable to underestimate host infection prevalence. Serological tests only document exposure to a pathogen, not infection. The tests for tick infection status are thought to be better, al-

though to our knowledge, no published information exists on sensitivity or specificity of any of the several commonly used diagnostic tests of *B. burgdorferi* in ticks—dark field microscopy (Junttila et al. 1999, Richter et al. 2000), direct fluorescent antibody assays (Burkot et al. 1994), and diagnostic and quantitative polymerase chain reaction (PCR) targeting the flagellin gene (Johnson et al. 1992, Schmidt et al. 1996, Junttila et al. 1999, Zeidner et al. 2001, Soares et al. 2006), the *recA* gene (Mommert et al. 2001, Wang et al. 2003), the *p66* gene (Mommert et al. 2001), the outer surface protein A gene (Persing et al. 1990), and several targets among the ribosomal genes (Kurtenbach et al. 1998, Kurtenbach et al. 2002, Layfield and Guilfoile 2002, Courtney et al. 2004, Schulze et al. 2005, Ornstein and Barbour 2006). Although several methods (e.g., quantitative real time PCR; Courtney et al. 2004, Ornstein and Barbour 2006) have been shown to detect vanishingly small amounts of bacteria from culture, it is not clear that this analytic sensitivity translates into the ability to correctly identify infected ticks (sensitivity in the epidemiological sense). Moreover, without a “gold standard” for *B. burgdorferi* infection, it has been impossible to estimate rates of false positives (imperfect specificity). Even a few false negative or false positive host test results could negatively or positively bias estimates of prevalence, particularly when sample sizes are low because of difficulties sampling rare and/or trap-shy species. Similarly, falsely negative or positive ticks would tend to depress or inflate estimates of infectivity. Thus, in addition to the methodological difficulties producing ecologically relevant measures of host prevalence and infectivity, it seems

Table 1. Values of prevalence (π) and infectivity (ϕ) for ten species of Northeastern vertebrates estimated from data by LoGiudice et al. 2003, either assuming the diagnostic test was perfect (no false negatives or false positives; left) or allowing for false positives (right)

Species	N [#]	Assuming perfect test			Allowing imperfect specificity (false positives)			Realized reservoir competence from LoGiudice et al.	
		Prevalence (π)		Infectivity (ϕ)	$\pi\phi^\S$	Prevalence (ϕ)		Infectivity (ϕ)	$\pi\phi$
		ML Estimate (SI) [°]	ML Estimate (SI)	ML Estimate (SI)		ML Estimate (SI)	N [#]		
White-footed mouse	35	1.0 (0.940–1.0)	0.914 (0.859–0.959)	0.914	1.0 (0.940–1.0)	0.911 (0.857–0.958)	0.911	0.921	27
Eastern chipmunk	51	0.971 (0.864–1.0)	0.569 (0.512–0.626)	0.552	0.868 (0.720–0.966)	0.659 (0.593–0.720)	0.572	0.550	43
White-tailed deer	19	0.311 (0.103–0.606)	0.150 (0.057–0.290)	0.047	0.088 (0.001–0.352)	0.222 (0.029–0.513)	0.020	0.046	5
Raccoon	13	1.0 (0.430–1.0)	0.017 (0.005–0.040)	0.017	0.0 (0.0–1.0)	0.257 (0.0–1.0)	NA	0.013	13
Virginia opossum	18	0.593 (0.261–0.990)	0.040 (0.018–0.081)	0.024	0.0 (0.0–1.0)	0.243 (0.0–0.200)	NA	0.026	19
Striped skunk	4	0.530 (0.106–0.945)	0.191 (0.096–0.318)	0.101	0.251 (0.013–0.738)	0.324 (0.149–0.526)	0.081	0.097	4
Short-tailed shrew	42	0.831 (0.690–0.934)	0.505 (0.455–0.555)	0.420	0.588 (0.423–0.740)	0.761 (0.700–0.816)	0.447	0.418	41
Masked shrew	11	0.701 (0.372–0.953)	0.537 (0.424–0.649)	0.377	0.528 (0.222–0.831)	0.627 (0.477–0.753)	0.331	0.512	7
Red & grey squirrel	2+7	0.831 (0.482–1.0)	0.061 (0.041–0.086)	0.051	0.279 (0.047–0.648)	0.081 (0.034–0.148)	0.023	0.147	2+7
					Specificity	0.968 (0.949–0.986)			

[°] Maximum likelihood estimate (support intervals—parameter values ± 2 log likelihood units from MLE).

[§] Realized reservoir competence is derived by multiplying π and ϕ .

[#] In some cases, our sample sizes were higher because we were able to use individual hosts from which few ticks were recovered, whereas LoGiudice et al. restricted their analyses to individuals from which 5 or more molted ticks were collected and tested.

inevitable that inaccurate diagnostic tests will bias these estimates.

Our purpose in this article is to provide a method for estimating the contributions of different host species to disease transmission or risk by separating the effects of prevalence and infectivity, and a means of accounting for imperfect diagnostic tests. We present a probabilistic model that uses infection data on fed ticks (or other vectors) from field-caught animals to decompose estimates of realized reservoir competence into their component parts: prevalence and infectivity (Fig. 1). We reanalyze data from LoGiudice et al. (2003) on the realized reservoir competence of 10 species (or groups of species in some cases) to estimate the relative contributions of prevalence and infectivity in determining ecological roles of host species in producing infected ticks. We then show how imperfect diagnostic tests of infection, particularly false positives, can bias metrics of reservoir competence, and then how false positive rates can be estimated from multispecies data to improve estimates of prevalence and infectivity.

Materials and Methods

Realized Reservoir Competence Data. LoGiudice et al. (2003) captured several ($n = 4$ –51) individuals of each of 10 vertebrate species (Table 1) from wild populations, brought them into the laboratory for 2–3 d to collect the ticks feeding on each animal after they had finished their bloodmeal. These ticks were allowed to molt and then screened for *B. burgdorferi* infection by using a direct immunofluorescent assay. Realized reservoir competence was then estimated as the number of positive ticks from an individual host divided by the number of ticks tested from that individual, averaged over all individuals of a species. If the number of ticks tested is the same for each animal, this amounts to dividing the number of positive ticks by the number of ticks

tested. As is often the case for vector burdens on individual hosts (e.g., Woolhouse et al. 1997), the number of ticks successfully recovered, molted, and tested from each animal varied considerably, and was often just one or two. LoGiudice et al. (2003) reasoned that individual hosts from which fewer than five ticks were recovered and tested would unacceptably bias realized reservoir competence estimates, so these individuals were excluded from analyses. In addition to this loss of data, the relative importance of prevalence and infectivity to these estimates cannot be resolved. The problem is that it is impossible to determine whether an individual host is infected when all of the ticks collected from it test negative. This problem, of course, becomes worse when there are fewer and fewer ticks collected from each individual. If we take a probabilistic approach, however, we can use all of the data available, even individuals from which a single tick is tested, and simultaneously estimate both prevalence and infectivity.

Statistical Analyses. We begin with the simplest case where we make the assumption that the test for *B. burgdorferi* is perfect, that is, it produces no false negatives or false positives (we relax this assumption below). We then simply need to understand the ways in which a tick can be infected by an individual host of a given species or remain uninfected (dashed box in Fig. 2). The probability that a single tick is positive is $\pi\phi$, where π is the probability the host is infected (prevalence) and ϕ is the probability that the individual host infects the tick, given that the host is infected (infectivity). There are two paths that can lead to an uninfected tick. The tick might feed on an infected host but remain uninfected, with probability $\pi(1 - \phi)$, or its host might be uninfected, which occurs at probability $(1 - \pi)$. The sum of these two paths is the probability of observing a single uninfected tick.

Often, however, many ticks feed on a host at the same time. Of these n feeding ticks, k of them are infected, where k can range from 0 to n . We can specify the probability that k of n ticks feeding on a given host




Fig. 2. The probability tree leading to a positive or negative tick (dashed inner box) and test result (entire figure). π is the probability the host is infected (i.e., prevalence), ϕ is the probability an infected host transmits the infection to the tick (infectivity), *Sensitivity* is the probability an infected tick tests positive (1 - false negative rate), and *Specificity* is the probability a negative vector tests negative (1 - false positive rate).

are infected in terms of prevalence (π) and infectivity (ϕ) with the binomial as

$$\pi \binom{n_i}{k_i} \phi^{k_i} (1 - \phi)^{n_i - k_i} + (1 - \pi) \binom{n_i}{k_i} 0^{k_i} 1^{n_i - k_i}.$$

If any of the n ticks are infected (i.e., $k \geq 1$), the term to the right of the plus sign collapses to zero because the host must be infected in order for the tick to have been infected. If none of the n ticks are infected ($k = 0$), then either the host was infected but did not transmit the infection, which is described by the term to the left of the plus sign, or it was uninfected, described by the term on the right. When $k = 0$ we cannot establish whether or not the host was infected and so the probabilities of both events must be taken into account.

Prevalence is a population-level parameter, specific to a given population of a particular host species. Infectivity, too, may be viewed as a population—if not species—specific constant, as discussed above. To estimate the population-level values of π and ϕ for a given species we simply find the values of each that maximize the likelihood of observing the k_i of n_i *B. burgdorferi*-infected ticks on each host, i , for $i = 1, 2, 3, \dots, N$, summed over all of the N hosts in the sample. The likelihood of the parameter estimates of π and ϕ given the data are

$$\mathcal{L}(\pi, \phi, | \text{data}) = \sum_{i=1}^N \left[\pi \binom{n_i}{k_i} \phi^{k_i} (1 - \phi)^{n_i - k_i} + (1 - \pi) \binom{n_i}{k_i} 0^{k_i} 1^{n_i - k_i} \right]. \quad [1]$$

Again, the second term on the right-hand side of equation 1 is zero if any of the ticks on host i are positive. In the appendix we provide code for finding the maximum likelihood estimates of π and ϕ using R (R Development Core Team 2005).

This method of estimating prevalence and infectivity from infection data for fed ticks makes two important assumptions. First, we assume all larval ticks feeding on hosts are uninfected. Previous studies have found little or no vertical transmission of *B. burgdorferi* (Piesman et al. 1986, Patrican 1997), supporting this assumption. Second, we assume that the diagnostic test is perfectly specific and sensitive, that is, there are no false positives or false negatives, respectively. If these values were known, they could be incorporated into the likelihood. As noted in the introduction, without a gold standard it is not possible to assess rates of false negatives and false positives. We therefore investigated the effects of false negative and false positive tests of tick infection status on estimates of prevalence and infectivity by simulating 1,000 “data sets” at each of nine levels of sensitivity (Fig. 3a and c) with specificity held constant at 1, and at nine levels of specificity (Fig. 3b and d), with sensitivity held constant at 1, for a total of 18,000 data sets. Each data set consisted of 20 hosts, each of which was randomly “infected” (or not) with probability = *Prevalence*. The 20 hosts were then assigned burdens of feeding vectors (V) by random draws from a Poisson distribution with mean = 10. The infection status of these feeding vectors was then determined by random draws from a binomial distribution with $n = V$ trials and the probability of success (i.e., infection) in each trial equal to *Infectivity* \times *Sensitivity* + (1 - *Infectivity*) (1 - *Specificity*) if the host was infected and (1 - *Specificity*) if the host was not. We then estimated the parameters π and ϕ for each data set by using equation 1 and compared these estimates with the “true” values. The importance of imperfect sensitivity and specificity was explored with several levels of true *Prevalence* and *Infectivity*, but we present only the results for *Prevalence* and *Infectivity* = 0.5.

Studies of realized reservoir competence often gather data from multiple host species at the same time. If we assume that false positives are a product of the diagnostic test rather than specific to a given host species, we can then use such multihost species data sets to estimate a common test specificity (=1 - rate of false positives). This assumption seems reasonable, particularly when the fed larvae are allowed to digest the host bloodmeal and molt, retaining little if any host-specific factors that might interfere with the diagnostic test, as was the case in the LoGiudice et al. (2003) data.

The logic to estimating specificity in addition to prevalence and infectivity is the same as in the simple case above except that now we 1) estimate species-specific prevalence and infectivity parameters π_j and ϕ_j from a data set of infection status of ticks fed on several different species; and 2) there is now a common diagnostic test-specific parameter,




Fig. 3. Violin plots showing the frequency of parameter estimates of prevalence (a and b) and infectivity (c and d) from simulated data sets with varying levels of test sensitivity and perfect specificity (a and c) and varying levels of test specificity with perfect sensitivity (b and d). Dashed gray lines (b and d) are the frequency of estimates when the actual level of test specificity was known. True prevalence and infectivity (thick horizontal lines) were both 0.5. At each level of specificity and sensitivity, prevalence and infectivity were estimated from each of 1,000 simulated data sets (data set consists of 20 hosts with an average of 10 vectors per host; details in Materials and Methods).

Sp , that allows for some false positives. The probability that a tick that fed on host species j (for species $j = 1, 2, 3, \dots, S$) becomes infected and tests positive is $\pi_j \phi_j$, again assuming the test is perfectly sensitive, although estimates are fairly robust to violations of this assumption (see Fig. 3). The probability that a tick that feeds on an infected individual of host species j , does not become infected, but still tests positive is $\pi_j (1 - \phi_j) (1 - Sp)$. Lastly, the probability that a tick feeds on an uninfected host of species j and still tests positive is $(1 - \pi_j) (1 - Sp)$. The likelihood becomes

$$\begin{aligned} \mathcal{L}(\pi_j, \phi_j, Sp \mid data) = & \sum_j \sum_i \left[\pi_j \binom{n_{ij}}{k_{ij}} [\phi_j + (1 - \phi_j)(1 - Sp)]^{k_{ij}} \right. \\ & \times [(1 - \phi_j) - (1 - \phi_j)(1 - Sp)]^{n_{ij} - k_{ij}} + (1 - \pi_j) \\ & \left. \times \binom{n_{ij}}{k_{ij}} [1 - Sp]^{k_{ij}} Sp^{n_{ij} - k_{ij}} \right]. \quad [2] \end{aligned}$$

Results

The maximum likelihood estimates of prevalence and infectivity, and their product, realized reservoir competence, vary considerably among species (Table 1). Some species tend not to be infected (e.g., white-tailed deer and striped skunks), whereas others are frequently infected (e.g., white-footed mice, eastern chipmunks). Species vary even more in terms of in-

fectivity. White-footed mice have a tremendous propensity to transmit *B. burgdorferi* to feeding ticks, whereas all others species have intermediate (e.g., eastern chipmunks and shrews) or low levels of infectivity (e.g., white-tailed deer and squirrels), which is in general agreement with previous studies (Donahue et al. 1987, Mather and Mather 1990, Mather et al. 1990, Telford et al. 1990, Norris et al. 1996, Ouellette et al. 1997, LoGiudice et al. 2003, Ullmann et al. 2003, but see Lane et al. 2005). The actual parameter estimates, however, can be strongly biased if we do not account for imperfections in the diagnostic test of tick infection status.

The simulation results demonstrated that false negatives are not as great a problem as false positives. Estimates of prevalence and infectivity were fairly robust to imperfectly sensitive diagnostic tests. The bulk of prevalence estimates clustered around the true value with as many as one in five positive vectors testing negative (*Sensitivity* = 0.8; Fig. 3a). (It is worth noting that the apparently wide range of prevalence estimates is a result of the simulation methods. Each simulated sample was drawn from a population with true prevalence = 0.5, but the actual prevalence within the sample varied. Simply scoring individual hosts as infected or not based on whether any ticks tested positive would have yielded similar levels of variability.) Infectivity became biased slightly low only when rates exceeded one in 40 false negatives (*Sensitivity* = 0.975; Fig. 3c).

These estimates are much more sensitive to imperfect specificity (false positives; Fig. 3b and d). Esti-

mated prevalence is biased high when the false positive rate exceeds ≈ 1 in 100 (*Specificity* = 0.99) and become wholly inaccurate by one in 20 (*Specificity* = 0.95). Essentially false positives cause uninfected hosts to seem positive—just one positive vector is evidence that the host it fed upon was infected. As such, the apparent prevalence is biased high. Conversely, infectivity is biased low with increasing rates of false positives. Falsely positive hosts do not, of course, transmit the (apparent) infection to many vectors, so only one or two of potentially many ticks test positive. These rare “transmission events” seem to be the result of low infectivity and thus diminish estimates of ϕ . In essence, it becomes impossible to distinguish whether a host species is frequently infected, but rarely transmits the infection to feeding vectors or whether vectors feeding on uninfected hosts falsely test positive. These biases are most pronounced when true prevalence is low and true infectivity is high (simulation results not shown). (In the case where false positive rates are higher than actual infectivity, the estimate of ϕ will instead be biased upward.)

Incorporating false positives into our analyses of the LoGiudice et al. (2003) data dramatically improved our model fit (Δ Akaike's Information Criterion adjusted for small sample size [AIC_c] = 250), supporting the conclusion that the direct immunofluorescent assay is not perfectly specific. We estimate specificity at ≈ 0.97 , meaning that three of 100 negative ticks are incorrectly scored as positive. Allowing for these false positives had the effect of reducing estimates of prevalence for almost all species, and estimates of infectivity generally increased. For two species, raccoons and opossums, the estimates are uninformative (Table 1). Individuals of these two species tended to have a single tick test positive; thus, they seemed to have high prevalence and low infectivity. With an imperfectly specific diagnostic test, however, these results may have instead been due to false positives. This suggests a limit to our ability to estimate prevalence and infectivity when realized reservoir competence converges on the estimated false positive rate ($1 - \text{Specificity}$). An experimental infection study of raccoons found that infectivity is, in fact, at or very near zero (Norris et al. 1996).

Discussion

Realized reservoir competence—the chance that a tick feeding on a particular free-ranging host species will become infected—is an ecologically and epidemiologically relevant metric. Species vary considerably in their realized reservoir competence (e.g., LoGiudice et al. 2003; Table 1), which in turn strongly influences the dynamics of and risk from *B. burgdorferi*. We can gain insights into why species vary in their ability to infect larval ticks by decomposing realized reservoir competence into prevalence and infectivity.

Mammal species have a wide-range of levels of *B. burgdorferi* prevalence (Table 1). This may reflect differences in exposure to infected nymphs (but see Mather et al. 1989) or species-specific differences in susceptibility to *B. burgdorferi* infection. Low infection

prevalence in some host species, such as white-tailed deer and striped skunks, that likely are repeatedly inoculated by infected nymphs (Magnarelli et al. 2004) suggests an innate immunity to *B. burgdorferi*. Immunity based on the complement pathway has been described for mule deer (*Odocoileus hemionus*) and western fence lizards (*Sceloporus occidentalis*) (Ullmann et al. 2003). What is clear is that not all species are equally infected. Nor are they equally infectious—we found considerable variability in infectivity. Infected white-footed mice are very likely to transmit *B. burgdorferi* to feeding larvae, whereas other hosts, such as squirrels, are generally dead-end hosts for the bacterium. Thus, even if infection prevalence is extremely high for all species, community composition will still be important to overall incidence of Lyme disease (LoGiudice et al. 2003, Keesing et al. 2006).

Interestingly, prevalence and infectivity are strongly positively correlated across the host species ($r = 0.88$, $t_{5df} = 4.1$, $P = 0.009$). This correlation suggests that both quantities are influenced by inherent physiological, likely immunological, features of each host, or by their rate of exposure to infected nymphs. Whatever keeps hosts infected and bacteremia high in the host—low rates of clearance by the immune system or high rates of (re)infection—will tend to increase the probability that the host transmits the infection to feeding larvae. Put another way, a cleared infection cannot be transmitted. The ability of infected white-footed mice to infect larval ticks decays with time, since infection (Donahue et al. 1987, Shih et al. 1995, Lindsay et al. 1997), although infections are not necessarily cleared (Hofmeister et al. 1999). The infectivity of meadow voles (Markowski et al. 1998, Anderson et al. 2006) and rice rats (Levin et al. 1995) also declines with time, suggesting a common pattern. Moreover, infectivity seems to differ between geographic locations (e.g., our estimate of $\phi \approx 0.9$ for white-footed mice in New York, compared with a maximal estimate of $\phi \approx 0.3$ in Ontario; Lindsay et al. 1997), in a way that is at least broadly consistent with differences in exposure to infected ticks. Thus, we would suggest that studies examining both the duration and dose-dependence of infectivity across species are likely to be fruitful and important.

We also must stress the importance of considering the accuracy of the diagnostic test. If we naively assumed that the direct fluorescent antibody (DFA) test correctly identified all negative ticks (i.e., no false positives), our estimates of prevalence would have been artificially high and our estimates of infectivity artificially low (Table 1). Indeed, when we allow for false positives we find that we cannot distinguish between low rates of transmission to ticks (low infectivity) and falsely positive test results for two species (i.e., raccoons and opossums). False positives are not unique to DFA, nor is the bias in parameter estimates unique to this likelihood-based method of estimation. Rather, these results demonstrate the limits of our knowledge when dealing with imperfectly specific tests. They also highlight an urgent need for data on the specificity of diagnostic tests. In the absence of such data, we urge scientists to be conservative when scoring infection status. It is encouraging to note, however, that if

the true test specificity is known and included in the likelihood calculations this bias is erased (Fig. 3b and d, dashed lines).

Last, we would like to emphasize that our reanalyses of the data of LoGiudice et al. (2003) strongly support their conclusions: different species have the potential to play dramatically different roles in *B. burgdorferi* transmission and Lyme disease risk. We find in general that those species most often infected are also the species with the greatest propensity to infect naïve larvae. It will be enlightening to see how widely this pattern holds in other vector-borne disease systems. The likelihood-based technique we presented should help researchers address this question.

Acknowledgments

We are grateful to Kelly Oggenfuss and a host of other field assistants for efforts in collecting data. The seeds of this article were planted during conversations with Mike Newhouse. Without Charles Canham's deep knowledge of all things statistical, that seed would never have developed and grown. This research was supported by grants from the National Science Foundation (most recently DEB 0444585), the National Institutes of Health (ROI AI053109), and the Andrew W. Mellon Foundation.

References Cited

- Anderson, J. M., K. I. Swanson, T. R. Schwartz, G. E. Glass, and D. E. Norris. 2006. Mammal diversity and infection prevalence in the maintenance of enzootic *Borrelia burgdorferi* along the western coastal plains of Maryland. *Vector Borne Zoonotic Dis.* 6: 411–422.
- Burkot, T. R., L. Patrican, and J. Piesman. 1994. Field trial of an outer surface protein A (OspA) antigen-capture enzyme-linked immunosorbent assay (ELISA) to detect *Borrelia burgdorferi* in *Ixodes scapularis*. *Am. J. Trop. Med. Hyg.* 50: 354–358.
- Courtney, J. W., L. M. Kostelnik, N. S. Zeidner, and R. F. Massung. 2004. Multiplex real-time PCR for detection of *Anaplasma phagocytophilum* and *Borrelia burgdorferi*. *J. Clin. Microbiol.* 42: 3164–3168.
- Donahue, J. G., J. Piesman, and A. Spielman. 1987. Reservoir competence of white-footed mice for Lyme disease spirochetes. *Am. J. Trop. Med. Hyg.* 36: 92–96.
- Eisendle, K., T. Grabner, and B. Zelger. 2007. Focus floating microscopy—"gold standard" for cutaneous borreliosis? *Am. J. Clin. Pathol.* 127: 213–222.
- Goffe, W. L., G. D. Ferrier, and J. Rogers. 1994. Global optimization of statistical functions with simulated annealing. *J. Econom.* 60: 65–99.
- Hofmeister, E. K., B. A. Ellis, G. E. Glass, and J. E. Childs. 1999. Longitudinal study of infection with *Borrelia burgdorferi* in a population of *Peromyscus leucopus* at a Lyme disease-zoonotic site in Maryland. *Am. J. Trop. Med. Hyg.* 60: 598–609.
- Johnson, B.J.B., C. M. Happ, L. W. Mayer, and J. Piesman. 1992. Detection of *Borrelia burgdorferi* in ticks by species specific amplification of the flagellin gene. *Am. J. Trop. Med. Hyg.* 47: 730–741.
- Junttila, T., M. Peltomaa, H. Soini, M. Marjamaki, and M. K. Viljanen. 1999. Prevalence of *Borrelia burgdorferi* in *Ixodes ricinus* ticks in urban recreational areas of Helsinki. *J. Clin. Microbiol.* 37: 1361–1365.
- Kahl, O., L. Gern, L. Eisen, and R. S. Lane. 2002. Ecological research on *Borrelia burgdorferi* sensu lato: terminology and some methodological pitfalls, pp. 29–46. In J. Gray, O. Kahl, R. S. Lane, and G. Stanek [eds.], *Lyme borreliosis: biology, epidemiology and control*. CABI, New York.
- Keesing, F., R. D. Holt, and R. S. Ostfeld. 2006. Effects of species diversity on disease risk. *Ecol. Lett.* 9: 485–498.
- Kurtenbach, K., M. Peacey, S.G.T. Rijpkema, A. N. Hoodless, P. A. Nuttall, and S. E. Randolph. 1998. Differential transmission of the genospecies of *Borrelia burgdorferi sensu lato* by game birds and small rodents in England. *Appl. Environ. Microbiol.* 64: 1169–1174.
- Kurtenbach, K., S. M. Schafer, H. S. Sewell, M. Peacey, A. Hoodless, P. A. Nuttall, and S. E. Randolph. 2002. Differential survival of Lyme borreliosis spirochetes in ticks that feed on birds. *Infect. Immun.* 70: 5893–5895.
- Lane, R. S., J. Mun, R. J. Eisen, and L. Eisen. 2005. Western gray squirrel (Rodentia: Sciuridae): a primary reservoir host of *Borrelia burgdorferi* in Californian oak woodlands? *J. Med. Entomol.* 42: 388–396.
- Layfield, D., and P. Guilfoile. 2002. The prevalence of *Borrelia burgdorferi* (Spirochaetales: Spirochaetaceae) and the agent of human granulocytic ehrlichiosis (Rickettsiaceae: Ehrlichiae) in *Ixodes scapularis* (Acari: Ixodidae) collected during 1998 and 1999 from Minnesota. *J. Med. Entomol.* 39: 218–220.
- Levin, M., J. F. Levine, C. S. Apperson, D. E. Norris, and P. B. Howard. 1995. Reservoir competence of the rice rat (Rodentia: Cricetidae) for *Borrelia burgdorferi*. *J. Med. Entomol.* 32: 138–142.
- Lindsay, L. R., I. K. Barker, G. A. Surgeoner, S. A. McEwen, and G. D. Campbell. 1997. Duration of *Borrelia burgdorferi* infectivity in white-footed mice for the tick vector *Ixodes scapularis* under laboratory and field conditions in Ontario. *J. Wildl. Dis.* 33: 766–775.
- LoGiudice, K., R. S. Ostfeld, K. A. Schmidt, and F. Keesing. 2003. The ecology of infectious disease: effects of host diversity and community composition on Lyme disease risk. *Proc. Natl. Acad. Sci. U.S.A.* 100: 567–571.
- Magnarelli, L. A., J. W. Ijdo, U. Ramakrishnan, D. W. Henderson, K. C. Stafford, and E. Fikrig. 2004. Use of recombinant antigens of *Borrelia burgdorferi* and *Anaplasma phagocytophilum* in enzyme-linked immunosorbent assays to detect antibodies in white-tailed deer. *J. Wildl. Dis.* 40: 249–258.
- Markowski, D., H. S. Ginsberg, K. E. Hyland, and R. J. Hu. 1998. Reservoir competence of the meadow vole (Rodentia: Cricetidae) for the Lyme disease spirochete *Borrelia burgdorferi*. *J. Med. Entomol.* 35: 804–808.
- Mather, T. N., and M. E. Mather. 1990. Intrinsic competence of three ixodid ticks (Acari) as vectors of the Lyme disease spirochete. *J. Med. Entomol.* 27: 646–650.
- Mather, T. N., S. R. Telford, S. I. Moore, and A. Spielman. 1990. *Borrelia burgdorferi* and *Babesia microti*: efficiency of transmission from reservoirs to vector ticks (*Ixodes dammini*). *Exp. Parasitol.* 70: 55–61.
- Mather, T. N., M. L. Wilson, S. I. Moore, J.M.C. Ribeiro, and A. Spielman. 1989. Comparing the relative potential of rodents as reservoirs of the Lyme disease spirochete (*Borrelia burgdorferi*). *Am. J. Epidemiol.* 130: 143–150.
- Mommert, S., R. Gutzmer, A. Kapp, and T. Werfel. 2001. Sensitive detection of *Borrelia burgdorferi sensu lato* DNA and differentiation of *Borrelia* species by LightCycler PCR. *J. Clin. Microbiol.* 39: 2663–2667.
- Norris, D. E., J. F. Levine, M. Menard, K. Nakagaki, P. Howard, and C. S. Apperson. 1996. Experimental infection of the raccoon (*Procyon lotor*) with *Borrelia burgdorferi*. *J. Wildl. Dis.* 32: 300–314.
- Ornstein, K., and A. G. Barbour. 2006. A reverse transcriptase-polymerase chain reaction assay of *Borrelia burgdorferi* 16S

- rRNA for highly sensitive quantification of pathogen load in a vector. *Vector Borne Zoonotic Dis.* 6: 103–112.
- Ostfeld, R., and F. Keesing. 2000. The function of biodiversity in the ecology of vector-borne zoonotic diseases. *Can. J. Zool.* 78: 2061–2078.
- Ouellette, J., C. S. Apperson, P. Howard, T. L. Evans, and J. F. Levine. 1997. Tick-raccoon associations and the potential for Lyme disease spirochete transmission in the coastal plain of North Carolina. *J. Wildl. Dis.* 33: 28–39.
- Patrican, L. A. 1997. Absence of Lyme disease spirochetes in larval progeny of naturally infected *Ixodes scapularis* (Acari: Ixodidae) fed on dogs. *J. Med. Entomol.* 34: 52–55.
- Persing, D. H., S. R. Telford, A. Spielman, and S. W. Barthold. 1990. Detection of *Borrelia burgdorferi* infection in *Ixodes dammini* ticks with the polymerase chain reaction. *J. Clin. Microbiol.* 28: 566–572.
- Piesman, J., J. G. Donahue, T. N. Mather, and A. Spielman. 1986. Transovarially acquired Lyme disease spirochetes (*Borrelia burgdorferi*) in field-collected larval *Ixodes dammini* (Acari: Ixodidae). *J. Med. Entomol.* 23: 219–219.
- R Development Core Team. 2005. R: a language and environment for statistical computing computer program, version 2.2.1. By R Development Core Team, Vienna, Austria.
- Richter, D., A. Spielman, N. Komar, and F. R. Matuschka. 2000. Competence of American robins as reservoir hosts for Lyme disease spirochetes. *Emerg. Infect. Dis.* 6: 133–138.
- Schauber, E. M., and R. S. Ostfeld. 2002. Modeling the effects of reservoir competence decay and demographic turnover in Lyme disease ecology. *Ecol. Appl.* 12: 1142–1162.
- Schmidt, B., R. R. Muellegger, C. Stockenhuber, H. P. Soyer, S. Hoedl, A. Luger, and H. Kerl. 1996. Detection of *Borrelia burgdorferi*-specific DNA in urine specimens from patients with erythema migrans before and after antibiotic therapy. *J. Clin. Microbiol.* 34: 1359–1363.
- Schulze, T. L., R. A. Jordan, C. J. Schulze, T. Mixson, and M. Papero. 2005. Relative encounter frequencies and prevalence of selected *Borrelia*, *Ehrlichia*, and *Anaplasma* infections in *Amblyomma americanum* and *Ixodes scapularis* (Acari: Ixodidae) ticks from central New Jersey. *J. Med. Entomol.* 42: 450–456.
- Shih, C. M., L. P. Liu, and A. Spielman. 1995. Differential spirochetal infectivities to vector ticks of mice chronically infected by the agent of Lyme disease. *J. Clin. Microbiol.* 33: 3164–3168.
- Soares, C.A.G., N. S. Zeidner, C. B. Beard, M. C. Dolan, G. Dietrich, and J. Piesman. 2006. Kinetics of *Borrelia burgdorferi* infection in larvae of refractory and competent tick vectors. *J. Med. Entomol.* 43: 61–67.
- Telford, S. R., T. N. Mather, G. H. Adler, and A. Spielman. 1990. Short-tailed shrews as reservoirs of the agents of Lyme disease and human babesiosis. *J. Parasitol.* 76: 681–683.
- Ullmann, A. J., R. S. Lane, K. Kurtenbach, M. Miller, M. E. Schrieffer, N. Zeidner, and J. Piesman. 2003. Bacteriolytic activity of selected vertebrate sera for *Borrelia burgdorferi* sensu stricto and *Borrelia bissettii*. *J. Parasitol.* 89: 1256–1257.
- Van Buskirk, J., and R. S. Ostfeld. 1995. Controlling Lyme disease by modifying the density and species composition of tick hosts. *Ecol. Appl.* 5: 1133–1140.
- Wang, G. Q., D. Liveris, B. Brei, H. Y. Wu, R. C. Falco, D. Fish, and I. Schwartz. 2003. Real-time PCR for simultaneous detection and quantification of *Borrelia burgdorferi* in field-collected *Ixodes scapularis* ticks from the northeastern United States. *Appl. Environ. Microbiol.* 69: 4561–4565.
- Wilske, B., V. Fingerle, and U. Schulte-Spechtel. 2007. Microbiological and serological diagnosis of Lyme borreliosis. *FEMS Immunol. Med. Microbiol.* 49: 13–21.
- Woolhouse, M.E.J., L. H. Taylor, and D. T. Haydon. 2001. Population biology of multihost pathogens. *Science (Wash., D.C.)* 292: 1109–1112.
- Woolhouse, M.E.J., C. Dye, J. F. Etard, T. Smith, J. D. Charlwood, G. P. Garnett, P. Hagan, J.L.K. Hii, P. D. Ndhlovu, R. J. Quinell, C. H. Watts, S. K. Chandiwana, and R. M. Anderson. 1997. Heterogeneities in the transmission of infectious agents: implications for the design of control programs. *Proc. Natl. Acad. Sci. U.S.A.* 94: 338–342.
- Zeidner, N. S., B. S. Schneider, M. C. Dolan, and J. Piesman. 2001. An analysis of spirochete load, strain, and pathology in a model of tick-transmitted Lyme borreliosis. *Vector Borne Zoonotic Dis.* 1: 35–44.

Received 14 May 2007; accepted 27 September 2007.

Appendix

R Code for Estimating Prevalence and Infectivity. We begin with the simple case, where there are no false positives. In this case we can estimate prevalence (π) and infectivity (ϕ) from data on a single species. The function `EstPrevInf()` calculates a vector of probabilities of observing `Pos`-infected vectors out of `Tested` vectors tested on each individual host, given $P = \pi$ and $I = \phi$.

```
EstPrevInf = function(P, I, Pos,
  Tested) {
  LL = log(P * dbinom(Pos, Tested,
    prob = I) + (1 - P) *
    dbinom(Pos, Tested, prob =
    0))
```

```
#to prevent the function from
#returning -inf, which causes
#errors in the optimization
#routine, those rows with -inf
```

`#are replaced with a very low value`

```
LL[is.infinite(LL)] = -99999
return(LL)
}
```

We then find the values of $P = \pi$ and $I = \phi$ that maximize the log likelihood of `EstPrevInf()`. For simple problems with few parameters, the `optim()` function of R is sufficient, but for more complex optimization problems we use a simulated annealing routine (Goffe et al. 1994) implemented in R (R Development Core Team 2005) by L. Murphy and C. Canham found in the Likelihood package at http://www.ecostudies.org/lme_R_code_tutorials.html. This routine, `anneal()`, also has the advantage of calculating support intervals on the parameter estimates. It sends a vector of predictions to a probability density function to calculate the probability of observing each. Because our predictions are actually probabilities, we use `PDF.wrapper()` to accept and send back these probabilities.


```
PDF.wrapper = function(predicted) {
  #just to keep anneal() happy
  return(predicted)
}
```

The `anneal()` function also requires a list of initial parameter estimates, `pars`, and lists of high (`par_hi`) and low (`par_lo`) boundaries for these parameters.

```
pars = list(P = 0.5, I = 0.5, Pos =
"Pos", Tested = "Tested", predicted =
"predicted")
par_lo = list(P = 0, I = 0)
par_hi = list(P = 1, I = 1)
```

The actual data (e.g., `ExampleData`) should be a data frame where each row refers to an individual animal and one column has the number of vectors Tested and another has number that test Pos.

```
ExampleData = data.frame(cbind(
  Tested = c(3, 10, 25, 16, 15, 12, 7,
  7, 3, 3, 2),
  Pos = c(3, 0, 13, 13, 8, 10, 7, 5,
  0, 0, 0)
))
```

We then call the simulated annealing optimization routine with the command:

```
Results = anneal(model = EstPrevInf,
par = pars, source_data =
  ExampleData, pdf = PDF.wrapper,
  par_lo, par_hi, dep_var =
  "Pos", initial_temp = 1
#lower temperature converges better
)
```

The parameter values of `Results$best_pars` should be $P = 0.661$ and $I = 0.689$, and the AIC_c , found in `Results$aic_corr`, should equal 48.48.

In the case where specificity is allowed to vary, we need to estimate its value across multiple species. The code is only slightly more complex, representing the π_j and ϕ_j parameters for each of the j species. Here, we present a simple example for just three species.

```
EstPrevInfSpec = function(I1, P1,
I2, P2, I3, P3, Spec, Pos, Tested,
Spp){
  LL = numeric(length(Pos))
  #First calculate the likelihoods
  #of the subset of individuals
  #that are species "One"
  LL[which(Spp == "One")] = log(
  #Positive tests from vectors fed on
  #infected animals
  P1*dbinom(Pos[which(Spp == "One")],
  Tested[which(Spp == "One")], prob =
  (I1 + (1 - I1)*(1 - Spec)))
  #Positives tests from vectors fed on
  #negative animals
  + (1 - P1)*dbinom(Pos[which(Spp ==
  "One")], Tested[which(Spp == "One")],
  prob = (1 - Spec))
  #Then for those that are species "Two"
  LL[which(Spp == "Two")] = log(P2*
  dbinom(Pos[which(Spp == "Two")],
```

```
Tested[which(Spp == "Two")], prob =
(I2 + (1 - I2)*(1 - Spec))) + (1-P2)
*dbinom(Pos[which(Spp == "Two")],
Tested[which(Spp == "Two")],
prob = (1 - Spec))
#Then finally those that are
#species "Three"
LL[which(Spp == "Three")] = log(P3*
dbinom(Pos[which(Spp == "Three")],
Tested[which(Spp == "Three")],
prob = (T3 + (1 - T3)*(1 - Spec))) +
(1 - P3)*dbinom(Pos[which(Spp ==
"Three")], Tested[which(Spp ==
"Three")], prob = (1 - Spec))
#to prevent the function from
#returning-inf, which causes
#errors in the optimization
#routine, those rows with -inf
#are replaced with a very low value
LL[is.na(LL)] = -99999
LL[is.infinite(LL)] = -99999
return(LL)
}
```

The parameter lists include more parameters, but are otherwise the same:

```
pars = list(I1 = 0.5, P1 = 0.5, I2 =
0.5, P2 = 0.5, I3 = 0.5, P3 = 0.5,
Spec = 0.5, Pos = "Pos," Tested =
"Tested," Spp = "Spp," predicted =
"predicted")
par_lo = list(I1 = 0, P1 = 0, I2 =
0, P2 = 0, I3 = 0, P3 = 0, Spec = 0)
par_hi = list(I1 = 1, P1 = 1, I2 =
1, P2 = 1, I3 = 1, P3 = 1, Spec = 1)
```

The data are also organized very similarly. There is just one additional column of data with the species names (here, species One, Two, and Three):

```
MultiSppData = data.frame(cbind(
  Tested = c(3, 10, 25, 16, 15, 12,
  7, 7, 3, 3, 2, 15, 16, 22, 25, 18,
  18, 16, 21, 23, 5, 9, 10, 12, 13,
  12, 8, 9, 17, 13, 6, 5, 16, 3, 1,
  1),
  Pos = c(3, 0, 13, 13, 8, 10, 7, 5,
  0, 0, 1, 5, 1, 6, 0, 2, 4, 3, 0, 0,
  0, 1, 2, 1, 1, 7, 6, 6, 7, 4, 3, 4,
  7, 2, 0, 1))
MultiSppData$Spp = c(rep("One", 11),
  rep("Two", 12), rep("Three", 13))
```

Last, the call to the `anneal()` function is almost identical, just specifying the `EstPrevInfSpec()` function and the new data, `MultiSppData`:

```
MultiSppResults = anneal(model =
  EstPrevInfSpec, par = pars, source_
  data = MultiSppData, pdf = PDF.
  wrapper, par_lo, par_hi, dep_var =
  "Pos", initial_temp = 1)
```

The resulting solution should have an AIC_c of 154.58, with parameters: $I1 = 0.674$, $P1 = 0.737$, $I2 = 0.191$, $P2 = 0.554$, $I3 = 0.505$, $P3 = 0.827$, and $Spec = 0.970$.