




	<i>Red Winged Blackbird</i>	<i>Marsh Wren</i>	<i>Swamp Sparrow</i>	<i>Yellow Warbler</i>	<i>Common Yellowthroat</i>	<i>Virginia Rail</i>	<i>Canada Goose</i>	<i>Least Bittern</i>	<i>American Goldfinch</i>	<i>Wood Duck</i>	<i>Song Sparrow</i>
<i>Iona</i>	77.12	0.98	0.33	2.94	0.65	4.58	5.23	0.33	1.63	0.33	1.31
<i>Constitution</i>	46.88	20.13	5.19	3.12	8.57	4.29	0.91	2.73	1.17	1.82	1.43
<i>Tivoli</i>	23.4	42.4	12.6	3.7	3.8	4.2	0.64	3.8	1.28	0.32	0.16
<i>Stockport</i>	33.6	27.1	11.2	8.9	6.8	3.7	0.93	1.87	2.34	-	2.57


Wells, A.W., Nieder WC, Swift BL, O'Connor KA, Weiss CA. (2008) Temporal changes in the breeding bird community at four Hudson River tidal marshes. J. of Coastal Research.





Daniels, R.A., K.E. Limburg, R.E. Schmidt, D.L. Strayer, and R.C. Chambers. 2005. Changes in Fish Assemblages in the Tidal Hudson River, New York. *American Fisheries Society Symposium*, 45:471-503.



Strayer, D. Long-term data on the freshwater tidal Hudson River.




Warren, Likens, Buso, & Kraft.
2008. Status and Distribution of
Fish in an Acid-Impacted
Watershed of the Northeastern
U.S. *Northeastern Naturalist*.



Incidence of Lyme Disease in the United States, 1991-2006. Source: www.textbookofbacteriology.net

Cases of Lyme Disease per 100,000 people in New York State Regions



Source: <http://lymebook.com>

