

Percentage of the eleven most common marsh-dependant bird species surveyed at each marsh in 2005.

	<i>Red Winged Blackbird</i>	<i>Marsh Wren</i>	<i>Swamp Sparrow</i>	<i>Yellow Warbler</i>	<i>Virginia Rail</i>	<i>Common Yellowthroat</i>	<i>Canada Goose</i>	<i>Least Bittern</i>	<i>American Goldfinch</i>	<i>Wood Duck</i>	<i>Song Sparrow</i>
<i>Iona</i>	77.12	0.98	0.33	2.94	0.65	4.58	5.23	0.33	1.63	0.33	1.31
<i>Constitution</i>	46.88	20.13	5.19	3.12	8.57	4.29	0.91	2.73	1.17	1.82	1.43
<i>Tivoli</i>	23.4	42.4	12.6	3.7	3.8	4.2	0.64	3.8	1.28	0.32	0.16
<i>Stockport</i>	33.6	27.1	11.2	8.9	6.8	3.7	0.93	1.87	2.34	-	2.57

Densities of marsh breeding bird nests (#nests / hectare)

	<i>Red Winged Blackbird</i>	<i>Marsh Wren</i>	<i>Virginia Rail</i>	<i>Swamp Sparrow</i>	<i>Gray Catbird</i>	<i>American Robin</i>	<i>Willow Flycatcher</i>	<i>American Goldfinch</i>	TOTAL
<i>Iona</i>	1.89	-	-	-	-	-	-	-	1.89
<i>Constitution</i>	3.77	13.68	-	-	-	-	-	-	17.45
<i>Tivoli</i>	0.49	60.49	1.46	0.98	-	-	1.46	-	64.88
<i>Stockport</i>	2.83	26.88	-	2.83	0.71	0.71	-	2.12	36.08

Wells, AW, Nieder WC, Swift BL, O'Connor KA, Weiss CA. (2008) Temporal changes in the breeding bird community at four Hudson River tidal marshes. *J. of Coastal Research*. SI 55 (221-235).