



Climate-dependent CO₂ emissions from lakes

Sarian Kosten,¹ Fábio Roland,² David M. L. Da Motta Marques,³ Egbert H. Van Nes,¹ Néstor Mazzeo,⁴ Leonel da S. L. Sternberg,⁵ Marten Scheffer,¹ and Jon J. Cole⁶

Received 24 June 2009; revised 20 December 2009; accepted 28 December 2009; published 8 May 2010.

[1] Inland waters, just as the world's oceans, play an important role in the global carbon cycle. While lakes and reservoirs typically emit CO₂, they also bury carbon in their sediment. The net CO₂ emission is largely the result of the decomposition or preservation of terrestrially supplied carbon. What regulates the balance between CO₂ emission and carbon burial is not known, but climate change and temperature have been hypothesized to influence both processes. We analyzed patterns in carbon dioxide partial pressure (*p*CO₂) in 83 shallow lakes over a large climatic gradient in South America and found a strong, positive correlation with temperature. The higher *p*CO₂ in warmer lakes may be caused by a higher, temperature-dependent mineralization of organic carbon. This pattern suggests that cool lakes may start to emit more CO₂ when they warm up because of climate change.

Citation: Kosten, S., F. Roland, D. M. L. Da Motta Marques, E. H. Van Nes, N. Mazzeo, L. da S. L. Sternberg, M. Scheffer, and J. J. Cole (2010), Climate-dependent CO₂ emissions from lakes, *Global Biogeochem. Cycles*, 24, GB2007, doi:10.1029/2009GB003618.

1. Introduction

[2] The importance of the world's oceans in global carbon cycling is well known and their influence on atmospheric CO₂ concentrations is explicitly incorporated in climate change models [*Intergovernmental Panel on Climate Change (IPCC)*, 2007]. So far, however, the role of inland waters has received less attention even though recent studies indicate that they play an important role in regulating carbon fluxes as well [*Cole et al.*, 2007; *Downing et al.*, 2008; *Duarte et al.*, 2008]. A significant part of the organic carbon initially sequestered as CO₂ by terrestrial ecosystems ends up in rivers and lakes. Only about half of this carbon is transported to the oceans [*Cole et al.*, 2007]. Much of the terrestrially produced carbon entering inland waters is buried in sediments or emitted as CO₂ to the atmosphere [*Cole et al.*, 2007]. In addition, primary production within inland waters represents a substantial carbon flux, especially in lakes with high concentrations of nutrients allowing high productivity [*Williamson et al.*, 2009]. This turns inland waters into

carbon processing hot spots in terrestrial landscapes and despite the fact that inland waters occupy a relatively small fraction of the Earth's surface, they play an important role in the global carbon cycle by processing large amounts of terrestrially derived carbon [*Battin et al.*, 2009]. Depending on the balance between processes such as respiration, primary production, groundwater carbon inflow and calcite precipitation, these systems may be carbon sinks, or become supersaturated with CO₂ and act as CO₂ sources to the atmosphere [*Cole et al.*, 1994, 2000; *Duarte and Prairie*, 2005; *Sobek et al.*, 2005]. All these processes are likely sensitive to changes in temperature and hydrology.

[3] Very little is known about the overall effects of climatic change on the carbon cycling in inland waters. Temperature, for example, may affect carbon cycles in a direct way through its influence on aquatic respiration [*Sand-Jensen et al.*, 2007] and primary production [*Flanagan et al.*, 2003], which may be most evident when it coincides with an increase in nutrient loading [*Christoffersen et al.*, 2006]. A differential temperature dependence of respiration and photosynthesis may lead to a decrease in carbon fixation and an increase in carbon emission [*Lopez-Urrutia et al.*, 2006]. Altered precipitation regimes may influence lakes' metabolism as well. For instance through its effect on the hydraulic residence time, which can have several effects including altering carbon sedimentation and mineralization [*Algesten et al.*, 2004; *Curtis*, 1998]; changing terrestrial inputs of nutrients and organic matter, and possibly primary production as well [*Reynolds*, 1994; *Schallenberg and Burns*, 1997]. Temperature and precipitation also have an indirect effect on lake's carbon cycle through their influence on terrestrial carbon fixation and the subsequent carbon leaching to the lake [*Sobek et al.*, 2005].

¹Department of Aquatic Ecology and Water Quality Management, Wageningen University, Wageningen, Netherlands.

²Laboratory of Aquatic Ecology, Institute of Biology, Federal University of Juiz de Fora, Juiz de Fora, Brazil.

³Instituto de Pesquisas Hidráulicas, Universidade Federal do Rio Grande do Sul, Porto Alegre, Brazil.

⁴Laboratorio de Ecología y rehabilitación de sistemas acuáticos, Facultad de Ciencias and CURE, Universidad de la República, Montevideo, Uruguay.

⁵Department of Biology, University of Miami, Coral Gables, Florida, USA.

⁶Cary Institute of Ecosystem Studies, Millbrook, New York, USA.




Figure 1. Location of the 82 South American lakes sampled.

[4] To explore the potential net effects of climate on carbon emissions from lakes, we sampled 82 comparable shallow lakes along a latitudinal gradient (5°–55°S) in South America (Figure 1) ranging in annual mean temperature from 4.0 to 27.6°C.

2. Methods

2.1. Site Description

[5] We sampled 82 lakes in the East of South America (Figure 1). Lakes were selected to resemble each other as much as possible morphologically, but to vary as much as possible in climate and, within climate regions, in trophic state. The lakes in our data set ranged from being oligotrophic to hypertrophic (Table 1). All lakes were shallow (maximum mean depth 4.5m) and relatively small (surface area ranged between 0.09 and 2.53 km²) (Table 1). The climate conditions at the sampling sites varied considerably; the most northern lake sites had maximum monthly air temperature up to 28.7°C whereas at the most southern lake locations the maximum was only 8.2°C [New *et al.*, 2002]. At the time of sampling the lake water temperature ranged from approximately 10 to 30°C. In each climate region, lakes were selected to vary as much as possible in trophic state (Table 1 and Kosten *et al.* [2009a]).

2.2. Sampling

[6] Lakes South of 25°S were sampled once during summer, the lakes nearer to the equator were sampled during dry season between November 2004 and March 2006.

[7] We collected water samples along the whole water column with a 1.5 m long tube at 20 random points in each lake between 0930 and 1200 LT. Two liters of each of these depth-integrated samples were gathered in a 40 L bucket resulting in a depth and area integrated bulk sample. Fil-

tration for various analyses in the laboratory was conducted directly after collection. Water and filters were then frozen until analysis. Sediment samples of the top 2 cm were taken in the center of the lake with a Kajak corer.

[8] Annual mean precipitation data were obtained from a high-resolution data set of surface climate over global land areas [New *et al.*, 2002].

2.3. Sample Analysis

[9] We determined the acid neutralizing capacity (ANC) titrimetrically using 0.05N HCl on unfiltered samples in the field directly after sampling. pH was determined also on unfiltered samples using a probe. Chlorophyll *a* (chl *a*) was extracted from filters (GF/C S&S) with 96% ethanol and absorbance was measured at 665 and 750 nm [Nusch, 1980]. For dissolved organic carbon, (DOC) analysis water was filtered through GF/F Whatman filters. DOC was determined by a Total Organic Carbon analyzer (Model 700, O.I.C International BV). As a measure for humic substances spectrophotometric light absorption at 380 nm was measured [Buiteveld, 1995] in filtered (0.45 μm S&S) water. The concentration of organic N and C in sediment was determined, after carbonate removal, using a CNS analyzer (NA-1500).

[10] The δ²H and δ¹⁸O of the bulk lake water sample were determined in a Multiflow system connected to an Isoprime Mass Spectrometer (Thermo Electron, Waltham, Massachusetts, United States). All isotopic ratios are expressed in δ units relative to the Vienna mean standard ocean water, here as

$$\delta D \text{ or } \delta^{18}O = \left[\frac{R_{\text{sample}}}{R_{\text{std}}} - 1 \right] \cdot 1000$$

in which R_{sample} and R_{std} are ²H/H or ¹⁸O/¹⁶O ratios of the sample and standard, respectively. The precision of analysis is ±2.0‰ and ±0.1‰ for δ²H and δ¹⁸O values, respectively.

2.4. Calculations and Data Analysis

[11] The pCO₂ was calculated from the pH and the ANC, adjusting for temperature, ionic strength and air pressure [Cole *et al.*, 1994]. Subsequently, the results were expressed as undersaturated or supersaturated with CO₂ relative to the atmosphere (RS). We used the global average atmospheric CO₂ pressure for the year 2005 as a reference (P. Tans,

Table 1. General Data on the Lakes Sampled

	Range	Mean	Median
Area (km ²)	0.09–2.53	0.62	0.46
Mean depth (m)	0.50–4.50	1.80	1.60
Conductivity (μS cm ⁻¹)	38–4930	527	167
Total nitrogen (mg N L ⁻¹)	<0.10–25.80	1.80	0.40
Total phosphorus (mg P L ⁻¹)	0.02–9.14	0.27	0.10
PVI (%) ^a	0–81	11	1
Chlorophyll <i>a</i> (μg L ⁻¹)	0.6–2889.0	79.6	4.7
The extinction of light at λ = 380 nm, as a proxy for humic substances (m ⁻¹)	0.3–54.2	5.9	4.0
Total suspended solids (mg L ⁻¹)	2–663	43	9
Light attenuation coefficient (m ⁻¹)	0.50–43.60	4.60	2.50

^aThe percentage of the lake's volume filled with submerged vegetation.




Figure 2. Frequency distribution of lakes undersaturated and supersaturated with CO₂ relative to the atmosphere. For undersaturation, the relative saturation (RS) = $-p\text{CO}_2(\text{air})/p\text{CO}_2(\text{water})$; for supersaturation, RS = $p\text{CO}_2(\text{water})/p\text{CO}_2(\text{air})$.

Trends in atmospheric carbon dioxide: Recent global CO₂, 2009, available at <http://www.esrl.noaa.gov/gmd/ccgg/trends/#global>. Lakes were classified as strongly CO₂ supersaturated (RS > 1.2); strongly undersaturated (RS < -1.2); or near saturation (-1.2 < RS < 1.2). The percentage of the lake's volume filled with submerged vegetation (PVI) was determined analogously to *Canfield et al.* [1984] (for details, see *Kosten et al.* [2009b]).

[12] As a proxy for the hydrological character of the lake we derived an inflow:evaporation ratio using $\delta^2\text{H}$ and $\delta^{18}\text{O}$ of the lake water. The derivation is based on the principle of light isotopes evaporating more quickly than heavy isotopes. The 'heavier' the lake water in comparison to the incoming water the more the lake water has been subject to evaporation. We calculated the inflow:evaporation ratio using the Gat-Bowser model [*Gat and Bowser*, 1991; *Rozanski et al.*, 2001]. The relative humidity input for the model was obtained from a global data set [*Bowen and Revenaugh*, 2003]. For the stable isotope composition of the inlet water we used the average composition of precipitation at the lake location as a proxy (extracted from map provided by G. J. Bowen (Waterisotopes.org, Purdue University, West Lafayette, 2003, available at http://wateriso.eas.purdue.edu/waterisotopes/pages/data_access/ArcGrids.html) on the basis of data from *Bowen and Revenaugh* [2003]). However, as the incoming water may have already been subject to evaporation in the watershed before it enters the lake, we may be overestimating the evaporation in the lake. The more the lake was subject to evaporation, for example due to a long hydraulic residence time, the smaller the inflow:evaporation ratio.

[13] Relations of $p\text{CO}_2$ versus climatological variables and local variables were first explored using simple linear regressions. The climatological variables included temperature, mean annual precipitation and the inlet:evaporation

ratio. The local variables included PVI, chlorophyll *a* (chl *a*), the extinction of light at $\lambda = 380$ nm used as a proxy for humic substances (humic), dissolved organic carbon concentration (DOC), and two indicators for the relative influence of aquatic primary production and terrestrial carbon input: (1) the ratio between chlorophyll *a* and the extinction at $\lambda = 380$ nm (chl *a*: humic); and (2) the ratio between organic carbon and nitrogen in the sediment (C:N). A high chl *a*: humic ratio indicates that the primary production is relatively large compared to influence of terrestrial organic matter on the lake's metabolism. To the contrary, a high C:N ratio indicates a relatively strong terrestrial influence on the lake's metabolism, as the C:N ratio of terrestrial organic matter is generally higher than that of aquatic material [*Elser et al.*, 2000]. Next, we applied multiple linear regressions. The multiple linear regressions modeled $p\text{CO}_2$ using temperature and the variables explaining most of the variance in $p\text{CO}_2$ in the simple linear regressions. To enhance normality, both the dependent and the independent (except for temperature) variables were log transformed before analysis.

[14] All statistical analyses were performed using SPSS for Windows version 15.0 (SPSS, Chicago, Illinois, United States).

3. Results

[15] Only in a small fraction (6%) of the lakes, the carbon dioxide partial pressure ($p\text{CO}_2$) was within a range of $\pm 20\%$ from the equilibrium with the atmosphere (near saturation group). Most lakes (80%) were supersaturated (relative saturation > 1.2, Figure 2). The $p\text{CO}_2$ saturation increased with increasing temperature (Table 2). Temperature alone explained 13% (R^2) of the variance in $p\text{CO}_2$ (Table 2). Lakes with high abundances of primary producers (either phytoplankton or submerged macrophytes) generally have lower $p\text{CO}_2$'s than expected based on temperature alone

Table 2. Simple Linear Regression Models Describing the Relationship of Lakes' Partial CO₂ Pressure and Different Climatological and Local Variables^a

Model	Log($p\text{CO}_2$) =	Regression Statistics		
		F ₁	R ²	n
1	$2.08^{(<0.001)} + 0.04^{(0.001)}$ temperature	12.13	0.13	82
2	$-0.17^{(0.877)} + 1.08^{(0.005)}$ log(total annual precipitation)	8.37	0.08	82
3	$2.55^{(<0.001)} + 0.98^{(<0.001)}$ log(inflow:evaporation) ^b	21.87	0.24	70
4	$3.33^{(<0.001)} - 0.30^{(<0.001)}$ log(chl <i>a</i>)	18.14	0.19	82
5	$3.15^{(<0.001)} - 0.18^{(0.047)}$ log(PVI) ^c	4.06	0.05	82
6	$3.10^{(<0.001)} - 0.08^{(0.548)}$ log(humic) ^d	0.36	0.01	82
7	$3.50^{(<0.001)} - 0.42^{(0.010)}$ log(DOC)	6.96	0.08	82
8	$3.20^{(<0.001)} - 0.36^{(0.001)}$ log(chl <i>a</i> : humic) ^e	21.63	0.21	82
9	$0.70^{(0.103)} + 2.27^{(0.001)}$ log(C:N) ^f	32.45	0.30	79

^aThe *p* value of the parameters are presented between brackets.

^bInflow:evaporation ratio.

^cThe percentage of the lake's volume filled with submerged vegetation (PVI) was augmented with 1% to avoid zero's.

^dThe extinction of light at $\lambda = 380$ nm is used as a proxy for humic substances.

^eChlorophyll *a* concentration:extinction at $\lambda = 380$ nm ratio.

^fOrganic carbon: total nitrogen ratio in the top sediment.




Figure 3. Partial CO₂ pressure ($p\text{CO}_2$) in lakes along a water temperature gradient in South America. Lakes differed in trophic status (size of symbols indicate chlorophyll a concentrations; open symbols represent lakes with substantial growth of submerged vegetation (>25% of the lake volume is filled with vegetation)). Continuous line represents the regression line between $p\text{CO}_2$ and the water temperature measured in the lakes. Lakes depicted below the dashed line are undersaturated in $p\text{CO}_2$ relative to the atmosphere.

(Figure 3, see also negative parameters in the models 4 and 5 in Table 2), whereas lakes with low abundances of primary producers generally fall above the $p\text{CO}_2$ temperature regression line.

[16] In the simple linear regressions $p\text{CO}_2$ was best explained by the inflow:evaporation ratio ($R^2 = 0.24$, model 3 in Table 2), the ratio between algal biomass (expressed as chl a) and the light extinction $\lambda = 380$ nm (used as a proxy for humic substances) ($R^2 = 0.21$, model 8 in Table 2), and the C:N ratio in the sediment ($R^2 = 0.30$, model 9 in Table 2). Lakes with a high inflow:evaporation ratio had relatively high $p\text{CO}_2$'s compared to lakes with a low inflow:evaporation ratio. Furthermore, in lakes with a relatively strong terrestrial influence (i.e., a low chl a : humic ratio or a high C:N ratio), the $p\text{CO}_2$ tended to be high as well.

[17] The $p\text{CO}_2$ could significantly be explained by a combination of temperature and the before mentioned variables, increasing the degree of explanation (adjusted R^2)

of $p\text{CO}_2$ (Table 3). At similar inflow:evaporation ratios, the cooler lakes had a lower $p\text{CO}_2$ than the warmer lakes (Table 3, Figure 4a). In the same way, the warm lakes generally had a higher $p\text{CO}_2$ than the cooler lakes at similar chl a : humic ratios (Table 3, Figure 4b).

4. Discussion

[18] The small fraction of lakes that is in equilibrium with the atmosphere indicates that most lakes were either sinks or sources to the atmosphere (Figure 2). As in surveys in other parts of the world [Cole *et al.*, 1994; Sobek *et al.*, 2003], most lakes were supersaturated, indicating that these lakes are net sources of CO₂ to the atmosphere. In fact, $p\text{CO}_2$ tends to be lowest in summer [Kelly *et al.*, 2001] or dry season [Richey *et al.*, 2002] precisely when our samples were taken. Therefore the annual average relative saturation per lake is likely even higher than we report here.

[19] The ratio between phytoplankton biomass and the light extinction at $\lambda = 380$ nm explained 21% (R^2) of the variation in $p\text{CO}_2$ (Table 2). Most likely this is because the light extinction at this wavelength, which we used as a proxy for the level of humic substances can be used as an indicator for terrestrial carbon input. When the terrestrial carbon is mineralized this may lead to a net CO₂ production. Phytoplankton, on the other hand, takes up CO₂ during photosynthesis, reducing CO₂ concentrations in the water. The higher the density of phytoplankton compared to the level of humic substances, the lower the $p\text{CO}_2$ tends to be. Besides phytoplankton, submerged vegetation may contribute substantially to the primary production in shallow lakes thereby lowering the $p\text{CO}_2$. Indeed, the $p\text{CO}_2$ is systematically lower in lakes with high abundances of submerged vegetation (Table 2 and Figure 3).

[20] In other lake data sets, $p\text{CO}_2$ was found to be positively correlated to DOC concentrations [Prairie *et al.*, 2002; Sobek *et al.*, 2003, 2005]. Although our lakes had a wide range in extinction at $\lambda = 380$ nm (used as a proxy for humic substances) and dissolved organic carbon (0.3–54.2 m^{-1} and 1.7–86 mg C L^{-1} , respectively), we did not find a significant relation between $p\text{CO}_2$ and the light extinction, and the explained variance in $p\text{CO}_2$ by DOC was low ($R^2 = 0.08$, Table 2). Our lakes, however, also covered a wide range in productivity and as $p\text{CO}_2$ is influenced by primary production and respiration simultaneously, the chlorophyll a :

Table 3. Multiple Linear Regression Models Describing the Relationship of Lakes' Partial CO₂ Pressure, Temperature, and Local Environmental Variables^a

		Coefficients ^b			Regression Statistics		
Intercept	Temperature	Inflow:evaporation	Chl a : humic	C:N		R^2_{adj}	n
1.93 ^(<0.001)	0.03 ^(0.035)	0.99 ^(<0.001)			$F_2 = 13.84$	0.27	70
2.20 ^(<0.001)	0.04 ^(<0.001)		-0.37 ^(<0.001)		$F_2 = 21.48$	0.34	82
2.17 ^(<0.0001)	0.03 ^(0.023)	0.67 ^(0.005)	-0.23 ^(0.011)		$F_3 = 12.29$	0.33	70
-0.03 ^(0.94)	0.04 ^(<0.001)			2.15 ^(<0.001)	$F_2 = 25.93$	0.39	79
0.48 ^(0.36)	0.03 ^(0.020)	0.54 ^(0.015)		1.62 ^(0.001)	$F_3 = 12.6$	0.35	67

^aIn the lower part we used the ratio between organic carbon and nitrogen in the sediment (C:N) instead of the chl a :humic ratio as an indicator for the relative influence of terrestrial and aquatic organic matter. Only significant models are shown. The p value of the parameters is presented between brackets. For explanation of the variables, see Table 2.

^bAll variables (except of temperature) were log transformed before analysis.




Figure 4. Partial CO₂ pressure ($p\text{CO}_2$) in South American lakes over a temperature gradient and a gradient of (a) a ratio between inflow and evaporation and (b) a ratio between algal biomass (expressed as chlorophyll *a*) and the extinction of light at $\lambda = 380$ nm, as a proxy for humic substances. The transparent planes indicate the multiple linear regression models. Grey points have a lower and black points have a higher $p\text{CO}_2$ than expected on the basis of the models.

humic substances ratio may therefore be most informative. A relatively constant primary production in comparison to the variation in DOC in the other data sets may explain the overruling influence of DOC on the $p\text{CO}_2$ in those earlier studies. The strong correlation between DOC and chlorophyll *a* in our data set ($R^2 = 0.48$, $p < 0.001$) might indicate that in eutrophic lakes, much of the DOC is algal derived which may explain the negative correlation between $p\text{CO}_2$ and DOC (Table 2).

[21] Our data furthermore suggest that hydrological characteristics of the lake, i.e., the ratio between inflow and evaporation, strongly affects the $p\text{CO}_2$ (Table 2). In part this may reflect a temperature effect. Evaporation plays an important role in the energy budget of the lake and strong evaporation cools down the lake [Lenters *et al.*, 2005]. Hydrology, however, also affects the input of terrestrial carbon to the lake. Indeed, $p\text{CO}_2$ was generally high in lakes where the volume of incoming water from the watershed was large in comparison to the volume that had evaporated. The importance of lakes' hydrology is in line with results from studies in other parts of the world [Algesten *et al.*, 2004]. Although the $p\text{CO}_2$ may be influenced by the inflow of CO₂ rich groundwater as well [Striegl and Michmerhuizen, 1998], a dominant role of the terrestrial input in our lakes is suggested by the fact that 21% of the variance in $p\text{CO}_2$ could be explained by the chlorophyll *a*: humic substances ratio.

[22] As an alternative indicator of the relative importance of terrestrial input, we analyzed the ratio between organic carbon and nitrogen in the sediment, a high C:N ratio indicating a relatively strong terrestrial influence on the lake's metabolism. Indeed we found this indicator to explain a comparable part (30%) of the variance in $p\text{CO}_2$ levels as the inflow:evaporation ratio (Table 2). Adding the ratios chlorophyll *a*: humic substances or C:N to regression models explaining $p\text{CO}_2$ levels from the hydrology indicator did not add to the explanatory power (Table 3), suggesting that the hydrology might affect the carbon balance largely

through its relative effects on allochthonous carbon input and in-lake primary productivity.

[23] For the lakes in our data set cooler lakes generally had a lower $p\text{CO}_2$ than warmer lakes (Table 3); the coldest lakes in our data set generally being carbon sinks and the warmer lakes in our data set being carbon sources (Figure 3). To filter out the effects of idiosyncratic differences in terrestrial inputs and productivity among lakes we looked at relationships between $p\text{CO}_2$ and the main indicators of these processes along the temperature gradient (Table 3, Figure 4). This again suggested a significant influence of temperature on the $p\text{CO}_2$. At similar inflow:evaporation ratios, or similar chlorophyll *a*: humic ratios the warmer lakes had a higher $p\text{CO}_2$ than the cooler lakes (Table 3, Figure 4). The logarithm of the $p\text{CO}_2$ in lakes with a water temperature of 30°C was 0.3 to 0.4 units higher than in lakes with a water temperature of 20°C (see temperature regression coefficients Table 3), suggesting an approximate 2–2.5 times increase in $p\text{CO}_2$ per 10°C in comparable lakes.

[24] Although the $p\text{CO}_2$ is not a direct measure of the carbon flux in lakes, it is the most important factor influencing this flux [Sobek *et al.*, 2005]. Therefore, our findings indicate a substantial influence of climate on CO₂ efflux from lakes. The apparent effect of hydrology on the $p\text{CO}_2$ in lakes suggested by our results implies that future changes in evaporation [Roderick and Farquhar, 2002] and precipitation regimes [IPCC, 2007] may have a strong impact on lake carbon emissions. In addition there is a clear relationship between $p\text{CO}_2$ and temperature. Although clearly correlations do not give insight in causal relationships, various mechanisms could contribute to such a temperature effect. In addition to the direct effect of temperature on the CO₂ flux (gasses dissolve better in colder than in warmer water), there is possibly an increase in net heterotrophy with temperature. Rates of respiration tend to increase stronger than production with temperature [Biddanda and Cotner, 2002; Lopez-Urrutia *et al.*, 2006; Rivkin and Legendre, 2001; Sand-Jensen *et al.*, 2007]. Importantly, the relatively strong

increase of respiration with temperature [Acuña *et al.*, 2008] implies that warm lakes might metabolize a substantially larger portion of the terrestrial organic matter influx than cooler ones [Biddanda and Cotner, 2002; Jansson *et al.*, 2008]. Primary production, on the other hand, may be limited by other factors than temperature (e.g., nutrients or light) this may prevent primary production from ‘keeping up’ with respiration when temperature increases.

5. Final Remarks

[25] Most likely, rising global temperatures will promote increased atmospheric greenhouse gas concentrations [Cox and Jones, 2008]. In fact, estimates based on ancient climate variation suggest that this effect may be quite large [Scheffer *et al.*, 2006]. Future climate prognosis often neglect the feedback effect of temperature on greenhouse gas fluxes as our knowledge of the processes involved is insufficient leading to large uncertainties [Jones *et al.*, 2006]. Our results suggest that warm inland waters emit more CO₂ than comparable cooler lakes. Carbon emission from cool lakes may therefore increase with climate warming. While this positive feedback mechanism has negative consequences from a climate change perspective, the sensitivity of the carbon balance of lakes to in-lake productivity and hydrology also suggests that a better understanding of the regulating mechanisms might give opportunities for managers to design climate friendlier management strategies of these hot spots of carbon channeling.

[26] **Acknowledgments.** We thank all people involved in the sampling and all lake owners that gave us access to the lakes. This research was financially supported by the Netherlands Organization for Scientific Research (NWO) grants W84-549 and WB840586, the National Geographic Society grant 7864-5; in Brazil by Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) grants 480122, 490409, and 311427; and in Uruguay by PEDECIBA, Maestría en Ciencias Ambientales, Donación Aguas de la Costa S.A., Banco de Seguros del Estado, and Sistema Nacional de Investigadores (SNI).

References

- Acuña, V., A. Wolf, U. Uehlinger, and K. Tockner (2008), Temperature dependence of stream benthic respiration in an Alpine river network under global warming, *Freshwater Biol.*, *53*, 2076–2088, doi:10.1111/j.1365-2427.2008.02028.x.
- Algesten, G., S. Sobek, A.-K. Bergstrom, A. Agren, L. J. Tranvik, and M. Jansson (2004), Role of lakes for organic carbon cycling in the boreal zone, *Global Change Biol.*, *10*, 141–147, doi:10.1111/j.1365-2486.2003.00721.x.
- Battin, T. J., S. Luyssaert, L. A. Kaplan, A. K. Aufdenkampe, A. Richter, and L. J. Tranvik (2009), The boundless carbon cycle, *Nat. Geosci.*, *2*, 598–600, doi:10.1038/ngeo618.
- Biddanda, B. A., and J. B. Cotner (2002), Love handles in aquatic ecosystems: The role of dissolved organic carbon drawdown, resuspended sediments, and terrigenous inputs in the carbon balance of Lake Michigan, *Ecosystems*, *5*, 431–445, doi:10.1007/s10021-002-0163-z.
- Bowen, G. J., and J. Revenaugh (2003), Interpolating the isotopic composition of modern meteoric precipitation, *Water Resour. Res.*, *39*(10), 1299, doi:10.1029/2003WR002086.
- Buiteveld, H. (1995), A model for calculation of diffuse light attenuation (PAR) and Secchi depth, *Neth. J. Aquat. Ecol.*, *29*, 55–65, doi:10.1007/BF02061789.
- Canfield, D. E., Jr., J. V. Shireman, D. E. Colle, W. T. Haller, C. E. Watkins, and M. J. Maccina (1984), Prediction of chlorophyll *a* concentrations in Florida lakes importance of aquatic macrophytes, *Can. J. Fish. Aquat. Sci.*, *41*, 497–501, doi:10.1139/f84-059.
- Christoffersen, K., N. Andersen, M. Søndergaard, L. Liboriussen, and E. Jeppesen (2006), Implications of climate-enforced temperature increases on freshwater pico- and nanoplankton populations studied in artificial ponds during 16 months, *Hydrobiologia*, *560*, 259–266, doi:10.1007/s10750-005-1221-2.
- Cole, J. J., N. F. Caraco, G. W. Kling, and T. K. Kratz (1994), Carbon dioxide supersaturation in the surface waters of lakes, *Science*, *265*, 1568–1570, doi:10.1126/science.265.5178.1568.
- Cole, J. J., M. L. Pace, S. R. Carpenter, and J. F. Kitchell (2000), Persistence of net heterotrophy in lakes during nutrient addition and food web manipulations, *Limnol. Oceanogr.*, *45*, 1718–1730.
- Cole, J. J., et al. (2007), Plumbing the global carbon cycle: Integrating inland waters into the terrestrial carbon budget, *Ecosystems*, *10*, 172–185, doi:10.1007/s10021-006-9013-8.
- Cox, P., and C. Jones (2008), Illuminating the modern dance of climate and CO₂, *Science*, *321*, 1642–1644, doi:10.1126/science.1158907.
- Curtis, P. (1998), Climatic and hydrological control of DOM concentration and quality in lakes, in *Aquatic Humic Substances*, edited by D. O. Hessen and L. J. Tranvik, pp. 93–105, Springer, Berlin.
- Downing, J. A., J. J. Cole, J. J. Middelburg, R. G. Striegl, C. M. Duarte, P. Kortelainen, Y. T. Prairie, and K. A. Laube (2008), Sediment organic carbon burial in agriculturally eutrophic impoundments over the last century, *Global Biogeochem. Cycles*, *22*, GB1018, doi:10.1029/2006GB002854.
- Duarte, C. M., and Y. T. Prairie (2005), Prevalence of heterotrophy and atmospheric CO₂ emissions from aquatic ecosystems, *Ecosystems*, *V8*, 862–870, doi:10.1007/s10021-005-0177-4.
- Duarte, C. M., Y. T. Prairie, C. Montes, J. J. Cole, R. Striegl, J. Melack, and J. A. Downing (2008), CO₂ emissions from saline lakes: A global estimate of a surprisingly large flux, *J. Geophys. Res.*, *113*, G04041, doi:10.1029/2007JG000637.
- Elsler, J. J., et al. (2000), Nutritional constraints in terrestrial and freshwater food webs, *Nature*, *408*, 578–580, doi:10.1038/35046058.
- Flanagan, K. M., E. McCauley, F. Wrona, and T. Prowse (2003), Climate change: The potential for latitudinal effects on algal biomass in aquatic ecosystems, *Can. J. Fish. Aquat. Sci.*, *60*, 635–639, doi:10.1139/f03-062.
- Gat, J. R., and C. Bowser (1991), The heavy isotope enrichment of water in coupled evaporative systems, in *Stable Isotope Geochemistry: A Tribute to Samuel Epstein*, edited by H. P. J. Taylor, J. R. O’Neil, and I. R. Kaplan, pp. 159–168, Geochem. Soc., St. Louis, Mo.
- Intergovernmental Panel on Climate Change (IPCC) (2007), *Climate Change 2007: Synthesis Report*, 52 pp., Intergov. Panel on Clim. Change, Geneva.
- Jansson, M., T. Hickler, A. Jonsson, and J. Karlsson (2008), Links between terrestrial primary production and bacterial production and respiration in lakes in a climate gradient in subarctic Sweden, *Ecosystems*, *11*, 367–376, doi:10.1007/s10021-008-9127-2.
- Jones, C. D., P. M. Cox, and C. Huntingford (2006), Climate-carbon cycle feedbacks under stabilization: Uncertainty and observational constraints, *Tellus, Ser. B*, *58*, 603–613.
- Kelly, C. A., E. Fee, P. S. Ramlal, J. W. M. Rudd, R. H. Hesslein, C. Anema, and E. U. Schindler (2001), Natural variability of carbon dioxide and net epilimnetic production in the surface waters of boreal lakes of different sizes, *Limnol. Oceanogr.*, *46*, 1054–1064.
- Kosten, S., V. L. M. Huszar, N. Mazzeo, M. Scheffer, L. da S. L. Sternberg, and E. Jeppesen (2009a), Lake and watershed characteristics rather than climate influence nutrient limitation in shallow lakes, *Ecol. Appl.*, *19*, 1791–1804, doi:10.1890/08-0906.1.
- Kosten, S., et al. (2009b), Climate-related differences in the dominance of submerged macrophytes in shallow lakes, *Global Change Biol.*, *15*, 2503–2517, doi:10.1111/j.1365-2486.2009.01969.x.
- Lenters, J. D., T. K. Kratz, and C. J. Bowser (2005), Effects of climate variability on lake evaporation: Results from a long-term energy budget study of Sparkling Lake, northern Wisconsin (USA), *J. Hydrol.*, *308*, 168–195, doi:10.1016/j.jhydrol.2004.10.028.
- Lopez-Urrutia, A., E. San Martin, R. P. Harris, and X. Irigoien (2006), Scaling the metabolic balance of the oceans, *Proc. Natl. Acad. Sci. U. S. A.*, *103*, 8739–8744, doi:10.1073/pnas.0601137103.
- New, M., D. Lister, M. Hulme, and I. Makin (2002), A high-resolution data set of surface climate over global land areas, *Clim. Res.*, *21*, 1–25, doi:10.3354/cr021001.
- Nusch, E. (1980), Comparison of different methods for chlorophyll and phaeopigments determination, *Arch. Hydrobiol.*, *14*, 14–36.

- Prairie, Y. T., D. F. Bird, and J. J. Cole (2002), The summer metabolic balance in the epilimnion of southeastern Quebec lakes, *Limnol. Oceanogr.*, *47*, 316–321.
- Reynolds, C. S. (1994), The ecological basis for the successful biomanipulation of aquatic communities, *Arch. Hydrobiol.*, *130*, 1–33.
- Richey, J. E., J. M. Melack, A. K. Aufdenkampe, V. M. Ballester, and L. L. Hess (2002), Outgassing from Amazonian rivers and wetlands as a large tropical source of atmospheric CO₂, *Nature*, *416*, 617–620, doi:10.1038/416617a.
- Rivkin, R. B., and L. Legendre (2001), Biogenic carbon cycling in the upper Ocean: Effects of microbial respiration, *Science*, *291*, 2398–2400, doi:10.1126/science.291.5512.2398.
- Roderick, M. L., and G. D. Farquhar (2002), The cause of decreased pan evaporation over the past 50 years, *Science*, *298*, 1410–1411.
- Rozanski, K., K. Froehlich, and W. G. Mook (2001), Surface water, in *Environmental Isotopes in the Hydrological Cycle: Principles and Applications*, edited by W. G. Mook, 121 pp., UNESCO/IAEA, Paris.
- Sand-Jensen, K. A. J., N. L. Pedersen, and M. Sondergaard (2007), Bacterial metabolism in small temperate streams under contemporary and future climates, *Freshwater Biol.*, *52*, 2340–2353, doi:10.1111/j.1365-2427.2007.01852.x.
- Schallenberg, M., and C. W. Burns (1997), Phytoplankton biomass and productivity in two oligotrophic lakes of short hydraulic residence time, *N. Z. J. Mar. Freshw. Res.*, *31*, 119–134.
- Scheffer, M., V. Brovkin, and P. M. Cox (2006), Positive feedback between global warming and atmospheric CO₂ concentration inferred from past climate change, *Geophys. Res. Lett.*, *33*, L10702, doi:10.1029/2005GL025044.
- Sobek, S., G. Algesten, A. N. N. K. Bergström, M. Jansson, and L. Tranvik (2003), The catchment and climate regulation of pCO₂ in boreal lakes, *Global Change Biol.*, *9*, 630–641, doi:10.1046/j.1365-2486.2003.00619.x.
- Sobek, S., L. Tranvik, and J. Cole (2005), Temperature independence of carbon dioxide supersaturation in global lakes, *Global Biogeochem. Cycles*, *19*, GB2003, doi:10.1029/2004GB002264.
- Striegl, R. G., and C. M. Michmerhuizen (1998), Hydrologic influence on methane and carbon dioxide dynamics at two north central Minnesota lakes, *Limnol. Oceanogr.*, *43*, 1519–1529.
- Williamson, C. E., J. E. Saros, and D. W. Schindler (2009), Sentinels of change, *Science*, *323*, 887–888, doi:10.1126/science.1169443.
-
- J. J. Cole, Cary Institute of Ecosystem Studies, Millbrook, NY 12545, USA.
- D. M. L. Da Motta Marques, Instituto de Pesquisas Hidráulicas, Universidade Federal do Rio Grande do Sul, Porto Alegre RS 91501-970, Brazil.
- S. Kosten, M. Scheffer, and E. H. Van Nes, Department of Aquatic Ecology and Water Quality Management, Wageningen University, PO Box 47, NL-6700 AA Wageningen, Netherlands. (sarian.kosten@wur.nl)
- N. Mazzeo, Laboratorio de Ecología y rehabilitación de sistemas acuáticos, Facultad de Ciencias and CURE, Universidad de la República, Iguá 4225, CP 11400 Montevideo, Uruguay.
- F. Roland, Laboratory of Aquatic Ecology, Institute of Biology, Federal University of Juiz de Fora, Juiz de Fora MG 36.036-900, Brazil.
- L. da S. L. Sternberg, Department of Biology, University of Miami, Coral Gables, FL 33124, USA.